



Airway obstruction and bronchial reactivity from age 1 month until 13 years in children with asthma

A prospective birth cohort study

Hallas, Henrik Wegener; Chawes, Bo Lund; Rasmussen, Morten Arendt; Arianto, Lambang; Stockholm, Jakob; Bønnelykke, Klaus; Bisgaard, Hans

Published in:
PLoS Medicine

DOI:
[10.1371/journal.pmed.1002722](https://doi.org/10.1371/journal.pmed.1002722)

Publication date:
2019

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Hallas, H. W., Chawes, B. L., Rasmussen, M. A., Arianto, L., Stockholm, J., Bønnelykke, K., & Bisgaard, H. (2019). Airway obstruction and bronchial reactivity from age 1 month until 13 years in children with asthma: A prospective birth cohort study. *PLoS Medicine*, 16(1), [e1002722]. <https://doi.org/10.1371/journal.pmed.1002722>

RESEARCH ARTICLE

Airway obstruction and bronchial reactivity from age 1 month until 13 years in children with asthma: A prospective birth cohort study

Henrik Wegener Hallas, Bo Lund Chawes[†], Morten Arendt Rasmussen[†], Lambang Arianto, Jakob Stokholm[†], Klaus Bønnelykke, Hans Bisgaard[†]*

Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte Hospital, University of Copenhagen, Copenhagen, Denmark

* bisgaard@copsac.com



Abstract

Background

Studies have shown that airway obstruction and increased bronchial reactivity are present in early life in children developing asthma, which challenges the dogma that airway inflammation leads to low lung function. Further studies are needed to explore whether low lung function and bronchial hyperreactivity are inherent traits increasing the risk of developing airway inflammation and asthmatic symptoms in order to establish timely primary preventive initiatives.

Methods and findings

We investigated 367 (89%) of the 411 children from the at-risk Copenhagen Prospective Studies on Asthma in Childhood 2000 (COPSAC₂₀₀₀) birth cohort born to mothers with asthma, who were assessed by spirometry and bronchial reactivity to methacholine from age 1 month, plethysmography and bronchial reversibility from age 3 years, cold dry air hyperventilation from age 4 years, and exercise challenge at age 7 years. The COPSAC pediatricians diagnosed and treated asthma based on symptom load, response to inhaled corticosteroid, and relapse after treatment withdrawal according to a standardized algorithm. Repeated measures mixed models were applied to analyze lung function trajectories in children with asthma ever or never at age 1 month to 13 years. The number of children ever versus never developing asthma in their first 13 years of life was 97 (27%) versus 270 (73%), respectively. Median age at diagnosis was 2.0 years (IQR 1.2–5.7), and median remission age was 6.2 years (IQR 4.2–7.8). Children with versus without asthma had reduced lung function (z-score difference, forced expiratory volume, -0.31 [95% CI -0.47 ; -0.15], $p < 0.001$), increased airway resistance (z-score difference, specific airway resistance, $+0.40$ [95% CI $+0.24$; $+0.56$], $p < 0.001$), increased bronchial reversibility (difference in change in forced expiratory volume in the first second [ΔFEV_1], $+3\%$ [95% CI $+2\%$; $+4\%$], $p < 0.001$), increased reactivity to methacholine (z-score difference for provocative dose, -0.40 [95% CI -0.58 ; -0.22], $p < 0.001$), decreased forced expiratory volume at cold dry air

OPEN ACCESS

Citation: Hallas HW, Chawes BL, Rasmussen MA, Arianto L, Stokholm J, Bønnelykke K, et al. (2019) Airway obstruction and bronchial reactivity from age 1 month until 13 years in children with asthma: A prospective birth cohort study. PLoS Med 16(1): e1002722. <https://doi.org/10.1371/journal.pmed.1002722>

Academic Editor: Aziz Sheikh, Edinburgh University, UNITED KINGDOM

Received: March 27, 2018

Accepted: November 30, 2018

Published: January 8, 2019

Copyright: © 2019 Hallas et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: COPSAC is funded by charitable and public research funds all listed on www.copsac.com. The Lundbeck Foundation (Grant no R16-A1694); The Ministry of Health (Grant no 903516); Danish Council for Strategic Research (Grant no 0603-00280B) and The Capital Region Research Foundation have provided core support for

COPSAC. The funding agencies did not have any role in design and conduct of the study; collection, management, and interpretation of the data; or preparation, review, or approval of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: COPSAC₂₀₀₀, Copenhagen Prospective Studies on Asthma in Childhood 2000; FEF₅₀, forced expiratory flow at 50% of expiration; FEV_{0.5}, forced expiratory volume in the first 0.5 seconds; FEV₁, forced expiratory volume in the first second; ICS, inhaled corticosteroid; MMEF, maximal mid-expiratory flow; PD₁₅, provocative dose of methacholine producing a 15% decrease in transcutaneous oxygen pressure; PD₂₀, provocative dose of methacholine producing a 20% decrease in forced expiratory volume in the first second; PtcO₂, transcutaneous oxygen pressure; sRaw, specific airway resistance.

challenge (Δ FEV₁, -4% [95% CI -7%; -1%], $p < 0.01$), and decreased forced expiratory volume after exercise (Δ FEV₁, -4% [95% CI -7%; -1%], $p = 0.02$). Both airway obstruction and bronchial hyperreactivity were present before symptom debut, independent of disease duration, and did not improve with symptom remission. The generalizability of these findings may be limited by the high-risk nature of the cohort (all mothers had a diagnosis of asthma), the modest study size, and limited ethnic variation.

Conclusions

Children with asthma at some point at age 1 month to 13 years had airway obstruction and bronchial hyperreactivity before symptom debut, which did not worsen with increased asthma symptom duration or attenuate with remission. This suggests that airway obstruction and bronchial hyperreactivity are stable traits of childhood asthma since neonatal life, implying that symptomatic disease may in part be a consequence of these traits but not their cause.

Author summary

Why was this study done?

- It is believed that asthma develops from inflammation in the lungs that leads to loss of lung function, but low lung function may be an inherent trait in children at risk of asthma instead of a consequence of inflammation.
- It is important to explore whether low lung function is an inherent trait that increases the risk of developing airway inflammation and asthma in order to establish primary preventive initiatives for low lung function.

What did the researchers do and find?

- Ninety-seven children who developed asthma and 270 children without asthma from the Danish COPSAC₂₀₀₀ birth cohort born to mothers with asthma were studied extensively with longitudinal lung function measurements from age 1 month to 13 years.
- Lung function was measured by spirometry from age 1 month and plethysmography from age 3 years, including assessments of bronchial reversibility to inhaled β_2 -agonist from age 3 years. Bronchial reactivity was assessed by methacholine challenge from age 1 month, cold dry air hyperventilation from age 4 years, and exercise challenge at age 7 years.
- Children developing asthma had reduced lung function from age 1 month throughout childhood compared to the children without asthma.
- The lung function deficit was present before the children developed asthma, did not progress with symptoms, and remained even if symptoms ceased.

What do these findings mean?

- Low lung function appears to begin in early childhood in a group of children who will develop asthma. The lung function trait is established prior to development of airway inflammation and asthma, and does not worsen with increased duration of asthma symptoms. As airway obstruction and increased bronchial reactivity manifest as early as 1 month of age, it is possible that preventive measures undertaken during pregnancy will be most effective.

Introduction

The current paradigm of asthma pathophysiology suggests that an inflammatory process in the airways leads to progressive bronchial hyperreactivity and lung function deficits [1]. An alternative hypothesis is causality in the opposite direction, i.e., that reduced airway caliber and bronchial hyperreactivity are inherent and stable traits that increase the risk of asthmatic symptoms, exaggerated hyperreactivity, and intermittent airway obstruction from a superimposed inflammatory process. In support of this, anti-inflammatory inhaled corticosteroid (ICS) treatment does not affect the natural course of lung function in children [2–6]. Furthermore, reticular basement membrane and airway smooth muscle thickness are unrelated to inflammatory cell counts in bronchial biopsies [7]. This alternative causal direction from airway obstruction and bronchial hyperreactivity to asthma symptoms implies that prevention must take place before symptom debut and maybe even before birth [8,9].

We aimed to study this alternative paradigm with rigorous prospective assessment of disease duration and remission. Spirometry and bronchial reactivity to methacholine were assessed in neonates [10,11]. Neonatal spirometry was volume-anchored and therefore comparable to repeated assessments of spirometry and bronchial reactivity during childhood [12–15]. Effort-independent measures of airway resistance and measures of bronchial reactivity were recorded until age 13 years.

Methods

This study is reported as per the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines ([S1 STROBE Checklist](#)).

The study was nested in Copenhagen Prospective Studies on Asthma in Childhood 2000 (COPSAC₂₀₀₀): a Danish prospective birth cohort study of 411 infants born during 1998–2001 to mothers with a history of asthma [16]. At enrollment at age 1 month, we excluded any child with a history of symptoms of lower airway infection or neonatal mechanical ventilation, pre-term birth (gestational age < 36 weeks), or any congenital abnormality or systemic illness. The children were examined at half-yearly scheduled visits until age 7 years, and again at age 13 years, including assessments of lung function 11 times during childhood. Thus, longitudinal lung function assessment during childhood was a strong and predefined focus of COPSAC₂₀₀₀, which is detailed in the cohort baseline paper [16], but there was no prospective analysis plan for how to model the data with respect to asthma development.

The local ethics committee (KF 01-289/96) and the Danish Data Protection Agency (2015-41-3696) approved the study. Both parents gave oral and written informed consent before enrollment.

Lung function

Neonatal spirometry. Neonatal spirometry was performed at age 1 month by forced flow volume measurements applying the volume-anchored raised volume rapid thoracoabdominal compression technique [10,11]. Forced expiratory volume in the first 0.5 seconds ($FEV_{0.5}$) and forced expiratory flow at 50% of expiration (FEF_{50}) were calibrated for gestational age at birth, sex, height, and weight.

Spirometry. Spirometry was performed half-yearly between age 5 and 7 years and at 13 years using a MasterScope Pneumoscreen spirometer (Erich Jaeger, Würzburg, Germany) [17]. Forced expiratory volume in the first second (FEV_1) and maximal mid-expiratory flow (MMEF) were calibrated for height, sex, and age.

To compare the neonatal $FEV_{0.5}$ and FEF_{50} with FEV_1 and MMEF data from later in childhood, the measurements were *z*-scored based on children never developing asthma and are subsequently referred to as FEV_z and $MMEF_z$.

Whole body plethysmography. Whole body plethysmography was performed half-yearly between age 3 and 7 years and at 13 years using a MasterScreen sealed bodybox (Erich Jaeger, Würzburg, Germany). Measurements were obtained during tidal breathing using a mouth-piece [18]. Specific airway resistance (sRaw) was calibrated for height, sex, and age; logarithmically transformed to achieve normal distribution of the data; and subsequently *z*-scored ($sRaw_z$).

Bronchial reversibility

Bronchial reversibility was measured at age 3, 5, 7, and 13 years by whole body plethysmography and at 5, 7, and 13 years by spirometry as the relative change from baseline 15 minutes after inhalation of β_2 -agonist. At age 3–7 years, terbutaline, 500 μ g ($2 \times 250 \mu$ g) pMDI (Bricanyl, AstraZeneca, Cambridge, UK) was delivered via a metal spacer (Nebuchamber, AstraZeneca, Cambridge, UK). At 13 years, we used salbutamol, 400 μ g ($2 \times 200 \mu$ g) from a discus DPI (Ventoline, GlaxoSmithKline Pharma, Middlesex, UK).

Bronchial reactivity

Methacholine challenge in neonates. Baseline $FEV_{0.5}$ was measured after a saline inhalation and following subsequent inhalations of methacholine in quadrupling dose steps from 0.04 to 16.67 μ mol/l delivered by a dosimeter attached to a nebulizer (SPIRA 08 TSM 133, Respiratory Care Center, Hämeenlinna, Finland) [10,11]. Bronchial reactivity was assessed from repeated measurements of $FEV_{0.5}$ and continuous transcutaneous oxygen pressure ($PtcO_2$) (TCM3, Radiometer, Copenhagen, Denmark) [10,11].

Methacholine challenge in school age children. Methacholine challenge in school-age children was performed at 7 and 13 years, measuring FEV_1 after a saline inhalation and after subsequent inhalations of methacholine in quadrupling dose steps from 0.06 to 29.72 μ mol/l (SPIRA 08 TSM 133, Respiratory Care Center, Hämeenlinna, Finland).

Bronchial reactivity to methacholine in neonates and school-age children was estimated from dose–response curves fitted with a logistic function as the provocative dose of methacholine producing a 15% decrease in $PtcO_2$ (PD_{15}) in the neonates, according to our previous sensitivity analysis in the cohort [11], and as the provocative dose of methacholine producing a 20% decrease in FEV_1 (PD_{20}) at age 7 and 13 years, according to the ATS/ERS guidelines [19]. To enable comparison of the assessments, PD_{15} and PD_{20} were logarithmically transformed to achieve normal distribution of the data, were *z*-scored, and are subsequently referred to as PD_z .

Cold dry air challenge. Cold dry air (-18°C) was delivered by a respiratory heat exchange system (Erich Jaeger, Würzburg, Germany) with an animated computer program guiding the child to hyperventilate at a level corresponding to 1 l/min/kg bodyweight for 4 minutes using a facemask with a mouthpiece [20]. Bronchial reactivity was determined as the change from baseline in sRaw at 4 and 6 years and in FEV₁ and MMEF at 6 years.

Exercise challenge. At 7 years, the child exercised on a motor-driven treadmill for at least 4 minutes with a pulse rate $>80\%$ of the maximal pulse [19] breathing fully dehumidified atmospheric air through a facemask (Hans Rudolph, Kansas City, MO, US). Spirometry was performed before, and 1, 3, 5, 10, and 15 minutes after exercise using the maximum percentage decline in FEV₁ from baseline within 10 minutes after exercise as the outcome.

We excluded lung function measurements in children with respiratory tract symptoms in the preceding week and if β_2 -agonists had been used within the preceding 12 hours. Bronchial challenge test results were excluded if the child had a respiratory infection in the preceding 3 weeks [21].

Monitoring of lung symptoms

The parents were taught to recognize troublesome lung symptoms including noisy breathing (wheezing or whistling sounds), breathlessness, shortness of breath, and persistent coughing at a session at the research unit. They subsequently recorded their child's troublesome lung symptoms, i.e., symptoms significantly affecting the well-being of their child, in a daily diary as a composite score (yes/no) from age 1 month to 7 years. They were requested to bring the child to the COPSAC clinic at each episode of 3 consecutive days with symptoms for examination and verification of the symptoms by the COPSAC pediatricians [2,22]. If the child was diagnosed with asthma, the diary recording of lung symptoms was continued beyond age 7.

Asthma diagnosis

Age 1 month–7 years. Asthma was diagnosed according to a predefined validated quantitative diary-based algorithm [22]. If the child experienced 5 episodes of troublesome lung symptoms within 6 months, daily symptoms for 4 weeks, or acute severe symptoms requiring hospitalization, high-dose ICS, or systemic corticosteroids, then a 3-month course of budesonide 200 μg bid (Spirocort, AstraZeneca, Cambridge, UK) was prescribed, and the child was examined by a chest X-ray and a sweat chloride test to exclude other chronic lung disorders. Asthma was diagnosed in children relapsing after ICS withdrawal within the following year, defined as either 2 episodes within 3 months or 2 consecutive weeks with symptoms, which resulted in a 6-month course of ICS and an additional 12-month course at subsequent relapses. Remission was defined as the child being without symptoms and treatment for 1 year.

After age 7 years. Children with troublesome lung symptoms after age 7 years were prescribed inhaled as-needed β_2 -agonist (terbutaline 250 μg ; Bricanyl, AstraZeneca, Cambridge, UK). If β_2 -agonist was needed more than 2 days per week and not only in relation to scheduled physical activity, the child was prescribed a 3-month course of ICS (budesonide 400 μg /day [$2 \times 200 \mu\text{g}$]; Spirocort, AstraZeneca, Cambridge, UK) and subsequently followed the same diagnostic procedure for asthma as described above. At the 13-year follow-up visit, the Danish National Medicine Registry was accessed and screened for any asthma drug prescriptions from age 7 to 13 years.

Only children attending the clinic at least once at age 0.5–4 years and at age 4–7 years or 13 years were included in the analyses.

Allergen sensitization

Allergen sensitization towards 10 aeroallergens—birch, grass, mugwort, horse, dog, cat, *Der-matophagoides pteronyssinus*, *D. farinae*, *Cladosporium herbarum*, and *Alternaria alternata*—was determined at age 13 years by skin prick test [23] and by measuring blood levels of specific IgE [24]. Allergic sensitization was defined as any wheal > 3 mm and/or any specific IgE value ≥ 0.35 kU/A.

Description of baseline characteristics of the cohort, including pre-, peri- and postnatal factors is provided in [S1 Text](#).

Statistical analyses

Baseline characteristics of included versus excluded children were compared using Fisher's exact test, Student's *t* test, and Wilcoxon 2-sample test.

The relationship between development of FEV_z, MMEF_z, sRaw_z, and PD_z in the first 13 years of life and asthma status was analyzed with mixed models taking repeated participant measurements into account. The statistic extracted from these mixed models was the fixed effect of group, i.e., asthma versus no asthma. To test the effect of the normalization of the lung function measurements, which was done to enable comparison of measurements conducted at varying ages, we also ran the analyses with raw, untransformed data and subsequently investigated the normal distribution of the residuals from the different models with Q-Q plots.

The cold dry air challenge by spirometry (6 years) and exercise challenge (7 years) were only performed once for each child, and the cross-sectional difference in these measures between children with asthma and without asthma was analyzed by Student's *t* test comparison of means.

We subsequently investigated whether development of FEV_z, MMEF_z, sRaw_z, and PD_z prior to onset of asthma was different from their development in children never developing asthma. This was done in mixed models comparing FEV_z, MMEF_z, sRaw_z, and PD_z measurements obtained before diagnosis in children developing asthma with measurements in children never developing asthma, evaluating the fixed effect of group, i.e., asthma versus no asthma. As an ancillary analysis, we also investigated by *t* tests whether neonatal FEV_{0.5}, FEV₅₀, and PD₁₅ were different in neonates who subsequently developed asthma versus neonates who did not develop asthma.

The effect of duration of asthma in the first 13 years of life on lung function development, i.e., whether lung function declined over time with increasing disease length, was investigated by calculating the duration of symptomatic disease preceding each lung function measurement. The mixed models used for this analysis were adjusted for time since remission and estimated the slope of the lung function during the period with asthma. The statistic extracted from these mixed models was the fixed effect of time, i.e., disease length in the observation period. To further investigate whether asthma status affected measures of FEV_z, MMEF_z, sRaw_z, and PD_z over time, age at measurement was included in the analyses as an interaction term.

Finally, lung function development after remission of asthma was analyzed in mixed models by evaluating the change over time in FEV_z, MMEF_z, sRaw_z, and PD_z after remission, evaluating the fixed effect of time since remission.

As post hoc analyses, we investigated development of FEV_z, MMEF_z, sRaw_z, and PD_z in children with remission of asthma, i.e., early-transient symptoms, compared to children with persistent symptoms and children never developing asthma. We also investigated development of lung function in children with asthma who had specific allergen sensitization at age 13 year

compared to children with asthma without sensitization. These post hoc analyses were done to investigate whether there were specific lung function trajectories in different wheeze phenotypes and whether development of atopy influenced the trajectories. The analyses were done with mixed models, evaluating the fixed effect of wheeze phenotype and sensitization status.

The effect of missing observations for longitudinal values for FEV_z , $MMEF_z$, $sRaw_z$, and PD_z was investigated by multiple imputation (for details see [S1 Text](#)).

All analyses were performed in R with the package *lme4* (version 1.1.14) [25] for linear mixed models as well as the package *mice* (version 3.3.0). All estimates are presented with 95% confidence intervals; p -values < 0.05 were considered significant. The graphical presentations of the results were made with *ggplot2* (version 3.0.0), and the longitudinal curves of FEV_z , $MMEF_z$, and $sRaw_z$ development were smoothed with loess regression function.

The data are available in [S1 Data](#).

Results

Of the 411 children enrolled at 1 month of age in COPSAC₂₀₀₀, follow-up to age 13 was available for 367 (89%). The number of children ever versus never developing asthma in their first 13 years of life was 97 (27%) versus 270 (73%), respectively ([Fig 1](#)). A total of 54 (56%) children remitted before age 13 years; 14 (14%) remitted but relapsed later during childhood. Median age at diagnosis was 2.0 years (IQR 1.2–5.7), median remission age was 6.2 years (IQR 4.2–7.8), and median duration of asthma was 4.0 years (IQR 2.2–5.3). The mean age of the mother at the birth of the child was 30.0 years, and 39% of the children had older siblings. There were no significant differences in baseline characteristics between included and excluded children ([Table 1](#)).

Lung function development in relation to asthma status

The development of FEV_z from age 1 month to 13 years in children ever versus never diagnosed with asthma in this period is depicted in [Fig 2A](#), illustrating that children developing asthma already had reduced FEV_z as neonates, which persisted until age 13 years as a stable trait without progression or attenuation during childhood (z -score difference, -0.31 [95% CI -0.47 ; -0.15], $p < 0.001$). The development of $MMEF_z$ was similar to that of FEV_z ([Fig 2B](#)) (z -score difference, -0.44 [95% CI -0.60 ; -0.27], $p < 0.001$). $sRaw_z$ measurements from age 3 to 13 years showed increased $sRaw$ from age 3 years among children ever versus never developing asthma, which was sustained till age 13 years without progression or attenuation ([Fig 2C](#)) (z -score difference, $+0.40$ [95% CI $+0.24$; $+0.56$], $p < 0.001$). Finally, bronchial reactivity to methacholine (PD_z) was persistently increased from age 1 month until age 13 years in children ever versus never diagnosed with asthma in that period ([Fig 2D](#)) (i.e., reduced PD_z ; z -score difference, -0.40 [95% CI -0.58 ; -0.22], $p < 0.001$) ([Table 2](#)). A sensitivity analysis utilizing the raw, untransformed data showed similar results ([Table A](#) in [S1 Text](#)). Furthermore, transforming the data led to more normalized residuals ([Fig A](#) in [S1 Text](#)). Finally, analyses using multiple imputation also showed the same differences in lung function development in children ever versus never diagnosed with asthma; although the effect estimates were slightly attenuated, the results remained significant ([Table B](#) in [S1 Text](#)).

In order to test whether the relationship between lung function development and asthma status varied with age, we introduced the interaction term asthma status \times age at measurement in the models. This analysis showed that there was no interaction between asthma status and age for lung function measurement ([Table C](#) in [S1 Text](#)), suggesting that these are predetermined traits established in early life and stable throughout childhood. An ancillary analysis of neonatal $FEV_{0.5}$, FEF_{50} , and PD_{15} showed significantly decreased forced flows and increased




Fig 1. Flowchart of included children.

<https://doi.org/10.1371/journal.pmed.1002722.g001>

Table 1. Baseline characteristics.

Characteristic	Included children (N = 367)	Excluded children (N = 44)	p-Value
Prenatal factors			
Boys, n/N (%) [*]	182/367 (50%)	21/44 (48%)	0.87
17q21 risk genotype ¹ , n/N (%) [*]	103/361 (29%)	11/32 (34%)	0.54
CDHR3 risk genotype ² , n/N (%) [*]	106/337 (31%)	7/28 (25%)	0.53
Filaggrin risk genotype ³ , n/N (%) [*]	44/363 (12%)	5/32 (16%)	0.57
Paternal asthma, n/N (%) [*]	54/358 (15%)	3/43 (7%)	0.17
Maternal smoking in third trimester, n/N (%) [*]	53/367 (14%)	10/44 (23%)	0.18
Perinatal factors			
Gestational age, mean (IQR) ^{**}	40 weeks (39; 41)	40 weeks (39; 41)	0.71
Maternal age at birth, mean (IQR) ^{**}	30.0 years (26.7; 33.2)	29.8 years (26.1; 33.7)	0.79
Cesarean section, n/N (%) [*]	80/367 (22%)	7/44 (16%)	0.44
Birth weight, mean (IQR) ^{**}	3.52 kg (3.20; 3.82)	3.52 kg (3.31; 3.79)	0.93
Birth length, mean (IQR) ^{**}	52.3 cm (51.0; 54.0)	52.4 cm (51.8; 53.3)	0.76
Postnatal factors			
Social circumstances, mean (SD) ^{**}	0.01 (1.00), N = 356	−0.19 (0.98), N = 26	0.30
Older sibling(s), n/N (%) [*]	139/358 (39%)	12/29 (41%)	0.84
Days of solely breastfeeding, median (IQR) ^{***}	122 (90; 155), N = 354	110 (14; 132), N = 13	0.26
Pets during first year, n/N (%) [*]	92/360 (26%)	12/35 (34%)	0.31
Age at start in daycare, median (IQR) ^{***}	12 months (8; 14), N = 367	13 months (8; 18), N = 34	0.33
Nicotine in hair at 3 years, GM (95% CI) ^{**}	0.37 ng (0.31; 0.44), N = 309	0.23 ng (0.08; 0.61), N = 11	0.31
Weight at 13 years, mean (IQR) ^{**}	49.2 kg (42.0; 55.1), N = 341	54.2 kg (47.0; 57.2), N = 4	0.46
Height at 13 years, mean (IQR) ^{**}	160.3 cm (154.3; 165.6), N = 341	162.4 cm (155.8; 165.9), N = 4	0.68

^{*}Fisher's exact test.

^{**}Student's *t* test.

^{***}Wilcoxon 2-sample test.

¹ Homozygous for the rs7216389 asthma risk allele (T).

² Hetero- or homozygous for the rs6967330 asthma risk allele (A).

³ Mutations in R501X, 2282del4, R2447X, and/or S3247X in the filaggrin coding gene.

GM, geometric mean.

<https://doi.org/10.1371/journal.pmed.1002722.t001>

reactivity to methacholine in neonates developing asthma within their first 13 years of life compared to neonates never diagnosed with asthma (Table D in S1 Text).

Children who developed asthma during the first 13 years of life compared to children never developing asthma also had increased bronchodilator response (absolute difference: FEV₁, +3% [95% CI +2%; +4%], *p* < 0.001; MMEF, +6% [95% CI 0%; 12%], *p* = 0.04; sRaw, −4% [95% CI −6%; −2%], *p* < 0.001), reduced post-bronchodilator *z*-scores for FEV₁ (−0.16 [95% CI −0.34; 0.01], *p* = 0.07) and MMEF (−0.35 [95% CI −0.54; −0.16], *p* < 0.001), and increased post-bronchodilator *z*-score for sRaw (+0.22 [95% CI +0.06; +0.38], *p* = 0.007).

Reactivity to cold dry air was increased among children ever versus never having asthma (absolute difference: FEV₁, −4% [95% CI −7%; −1%], *p* = 0.007; MMEF, −8% [95% CI −15%; −1%], *p* = 0.03; and sRaw, +9% [95% CI +5%; +13%], *p* < 0.001). Additionally, children ever versus never diagnosed with asthma experienced a significantly larger drop in FEV₁ after exercise challenge at age 7 years (absolute difference: FEV₁, −4% [95% CI −7%; −1%], *p* = 0.02) (Table 2).




Fig 2. Development of z-scored measures of lung function and bronchial reactivity from 1 month to 13 years. z-Scores for (A) forced expiratory volume (FEV_z), (B) maximal mid-expiratory flow (MMEF_z), (C) specific airway resistance (sRaw_z), and (D) provocative dose of methacholine (PD_z). Dashed lines represent 95% confidence intervals; grey triangles indicate timing of measurements. The vertical bars in (D) indicate the exact distribution in terms of age of sampling of the raw data.

<https://doi.org/10.1371/journal.pmed.1002722.g002>

Lung function development in relation to debut, duration, and remission of asthma

Children who developed asthma showed significantly increased airway obstruction and bronchial reactivity before symptom debut compared to children never developing asthma (FEV_z,

Table 2. Lung function and bronchial reactivity in relation to asthma at age 1 month to 13 years.

Measure	Asthma	N	Mean	Difference (95% CI)	p-Value
FEV					
FEV _z z-score [#]	Never	270	0.00	Reference	<0.001
	Ever	97	-0.33	-0.31 (-0.47; -0.15)	
FEV _z z-score after β ₂ -agonist [#]	Never	261	0.00	Reference	0.07
	Ever	94	-0.20	-0.16 (-0.34; +0.01)	
Percent change in FEV ₁ after β ₂ -agonist [#]	Never	261	+3%	Reference	<0.001
	Ever	94	+6%	+3% (+2%; +4%)	
Percent change in FEV ₁ after cold dry air ^{##}	Never	182	-3%	Reference	0.007
	Ever	69	-7%	-4% (-7%; -1%)	
Percent change in FEV ₁ after exercise ^{##}	Never	210	-8%	Reference	0.02
	Ever	78	-12%	-4% (-7%; -1%)	
MMEF					
MMEF _z z-score [#]	Never	269	0.00	Reference	<0.001
	Ever	97	-0.46	-0.44 (-0.60; -0.27)	
MMEF _z z-score after β ₂ -agonist [#]	Never	261	0.00	Reference	<0.001
	Ever	94	-0.38	-0.35 (-0.54; -0.16)	
Percent change in MMEF after β ₂ -agonist [#]	Never	261	+21%	Reference	0.04
	Ever	94	+28%	+6% (0%; +12%)	
Percent change in MMEF after cold dry air ^{##}	Never	178	-4%	Reference	0.03
	Ever	68	-12%	-8% (-15%; -1%)	
sRaw					
sRaw _z z-score [#]	Never	263	0.00	Reference	<0.001
	Ever	94	+0.41	+0.40 (+0.24; +0.56)	
sRaw _z z-score after β ₂ -agonist [#]	Never	261	0.00	Reference	0.007
	Ever	93	+0.25	+0.22 (+0.06; +0.38)	
Percent change in sRaw after β ₂ -agonist [#]	Never	261	-19%	Reference	<0.001
	Ever	93	-23%	-4% (-6%; -2%)	
Percent change in sRaw after cold dry air ^{##}	Never	215	+8%	Reference	<0.001
	Ever	76	+16%	+9% (+5%; +13%)	
PD					
PD _z z-score [#]	Never	267	0.00	Reference	<0.001
	Ever	95	-0.39	-0.40 (-0.58; -0.22)	

Significant p-values in bold.

[#]Mixed models.

^{##}Student's *t* test.

FEV, forced expiratory volume; MMEF, maximal mid-expiratory flow; PD, provocative dose of methacholine; sRaw, specific airway resistance.

<https://doi.org/10.1371/journal.pmed.1002722.t002>

−0.29 [95% CI −0.49; −0.09], $p = 0.004$; MMEF_z, −0.40 [95% CI −0.60; −0.20], $p < 0.001$; sRaw_z, +0.33 [95% CI +0.10; +0.57], $p = 0.006$; and PD_z, −0.35 [95% CI −0.59; −0.12], $p = 0.003$).

The airway obstruction and increased bronchial reactivity in children developing asthma were independent of disease duration (change per year: FEV_z, −0.01 [95% CI −0.07; +0.06], $p = 0.82$; MMEF_z, −0.01 [95% CI −0.05; +0.03], $p = 0.68$; sRaw_z, −0.02 [95% CI −0.06; +0.03], $p = 0.43$; and PD_z, −0.04 [95% CI −0.10; +0.03], $p = 0.27$). Furthermore, the airway obstruction and increased bronchial reactivity did not attenuate after remission (change per year: FEV_z, +0.02 [95% CI −0.03; +0.06], $p = 0.44$; MMEF_z, −0.00 [95% CI −0.04; +0.04], $p = 0.98$; sRaw_z, +0.01 [95% CI −0.03; +0.06], $p = 0.54$; and PD_z, +0.00 [95% CI −0.07; +0.08], $p = 0.92$) (Table 3).

Lung function development in relation to transient versus persistent asthmatic symptoms

Among the 97 children who fulfilled the diagnostic criteria for asthma, 54 children had a transient phenotype and remitted before age 13 years, whereas the remaining 43 children had persistent asthmatic symptoms requiring continued ICS by age 13 years. Lung function development in children with transient asthmatic symptoms compared to children never diagnosed with asthma showed reduced FEV_z (z-score difference, −0.30 [95% CI −0.51; −0.08], $p = 0.008$) and reduced MMEF_z (−0.36 [95% CI −0.59; −0.14], $p = 0.002$) from age 1 month to 13 years, and increased sRaw_z from age 3 years to 13 years (+0.27 [95% CI +0.06; +0.48], $p = 0.01$). PD_z was reduced among transiently asthmatic versus healthy children, i.e., indicating increased airway reactivity, but this difference was not significant (−0.15 [95% CI −0.39; +0.09], $p = 0.23$).

There were no differences in development of FEV_z, MMEF_z, or sRaw_z among children with transient versus persistent asthmatic symptoms, whereas children with transient compared to

Table 3. Lung function development in relation to asthma debut, duration, and remission.

Measure	N	Difference (95% CI)	p-Value
z-Score before diagnosis relative to children never developing asthma			
FEV _z before diagnosis [#]	93	−0.29 (−0.49; −0.09)	0.004
MMEF _z before diagnosis [#]	93	−0.40 (−0.60; −0.20)	<0.001
sRaw _z before diagnosis [#]	32	+0.33 (+0.10; +0.57)	0.006
PD _z before diagnosis [#]	87	−0.35 (−0.59; −0.12)	0.003
Change in z-score per year duration of asthma			
FEV _z development [#]	92	+0.02 (−0.02; +0.07)	0.36
MMEF _z development [#]	92	−0.01 (−0.05; +0.03)	0.68
sRaw _z development [#]	89	−0.02 (−0.06; +0.03)	0.43
PD _z development [#]	87	−0.04 (−0.10; +0.03)	0.27
Change in z-score per year after remission of asthma			
FEV _z development [#]	54	+0.02 (−0.03; +0.06)	0.44
MMEF _z development [#]	54	−0.00 (−0.04; +0.04)	0.98
sRaw _z development [#]	51	+0.01 (−0.03; +0.06)	0.54
PD _z development [#]	48	+0.00 (−0.07; +0.08)	0.92

Significant p -values in bold.

[#]Mixed models.

FEV, forced expiratory volume; MMEF, maximal mid-expiratory flow; PD, provocative dose of methacholine; sRaw, specific airway resistance.

<https://doi.org/10.1371/journal.pmed.1002722.t003>

persistent symptoms had less airway reactivity to methacholine, i.e., higher PD_z (+0.48 [95% CI +0.15; +0.80], $p = 0.004$) (Fig 3; Table E in S1 Text).

Lung function development in relation to allergic sensitization

At age 13 years, 157 (52%) of 304 tested children had allergic sensitization towards 1 or more of the tested allergens by either a positive skin prick test and/or elevated blood level of specific IgE. Sensitization was diagnosed in 109 (49%) of 223 children never developing asthma, 15 (36%) of 42 children with transient asthmatic symptoms, and in 33 (85%) of 39 children with persistent asthmatic symptoms. Thus, sensitization was more prevalent in children with persistent asthma versus children never developing asthma (chi-squared test, $p < 0.001$) and in children with persistent versus transient asthmatic symptoms (chi-squared test, $p < 0.001$).

There were no differences in development of FEV_z , $MMEF_z$, or $sRaw_z$ among children who developed asthma with versus without allergen-specific sensitization to any of the tested allergens, whereas there was a pattern suggesting increased airway reactivity to methacholine among children with asthma who were sensitized by age 13 years, i.e., lower PD_z (−0.34 [95% CI −0.70; +0.02], $p = 0.06$) (Table E in S1 Text). Stratifying the analyses on type of allergen as (1) house dust mite, (2) furred animal (dog, cat, horse), or (3) pollen (birch, grass, mugwort) showed similar results except that children with asthma and sensitization to pollen had significantly lower PD_z (Table E in S1 Text).

Discussion

Primary findings

Children developing asthma at any time during the first 13 years of life had airway obstruction and increased bronchial reactivity that was present at age 1 month without worsening with increased symptom duration and without improvement after remission. This suggests that these are inherent and stable traits not caused by inflammation during symptomatic periods but rather that predispose the child to develop asthmatic symptoms, exaggerated hyperreactivity, and intermittent airway obstruction.

Strengths and limitations

The primary strength of this study is the thorough single-center close clinical surveillance of a cohort with a follow-up of 89% till age 13 years. The children were seen for clinical evaluation and lung function assessment at the age of 1 month and thereafter at least half-yearly until age 7 years and again at 13 years. Until age 7 years, the parents filled daily diary cards describing the child's troublesome lung symptoms, thereby diminishing recall bias [26]. Asthma was diagnosed and treated by the COPSAC pediatricians only according to a predetermined algorithm with a conservative definition based on symptom load and response to and relapse after a standard dose and period of ICS [22], thus avoiding diagnostic heterogeneity. Children with asthma were seen for scheduled 3-monthly visits and for unscheduled acute visits during worsening of symptoms and continued daily symptom diary recording beyond age 7 years if symptoms persisted.

The diagnostic algorithm for asthma, which was based on quantitative symptom assessments verified at scheduled and acute care visits to the research unit and response to ICS treatment, could identify asthmatic children with transient, recurrent viral-induced wheeze and children with a persistent phenotype with or without atopy. This is apparent as approximately half of the children diagnosed according to the algorithm had transient symptoms and remitted before age 13 years. The median age at diagnosis was 2 years, implying that a large




Fig 3. Development of z-scored measures of lung function and bronchial reactivity from 1 month to 13 years in children with transient asthma, children with persistent asthma, and children never developing asthma. z-Scores for (A) forced expiratory volume (FEV_z), (B) maximal mid-expiratory flow ($MMEF_z$), (C) specific airway resistance ($sRaw_z$), and (D) provocative dose of methacholine (PD_z). Dashed lines represent 95% confidence intervals; grey triangles indicate timing of measurements.

<https://doi.org/10.1371/journal.pmed.1002722.g003>

proportion of the cases were children whose initial episode of wheezing was related to an infection, but half of these children had ICS-dependent asthma by age 13 years. Importantly, not all children with asthma will respond to ICS initially, and because lack of ICS response cannot exclude asthma, we excluded other chronic lung diseases by means of chest X-ray and a sweat chloride test for cystic fibrosis.

A significant advantage of the study is the longitudinal repetitive lung function assessments performed at 11 time points during childhood by means of spirometry and plethysmography as well as bronchial reactivity to cold dry air, exercise, and methacholine challenge in accordance with recognized guidelines [17–20,27]. We did most measurements in the age range from 3 to 7 years; therefore, the shape of the curves is likely to be influenced by the readings at 1 month and 13 years. Still, the birth cohort design assured that lung function was assessed (1) before the debut of symptoms and asthma diagnosis, (2) repeatedly every 6 months in children with and without asthma, and (3) after remission in children outgrowing their asthma. This allowed us to scrutinize how lung function trajectories develop in children with asthma during childhood and how they are affected by duration and remission of symptoms. Our interaction analyses showed no evidence of age influence on the difference in lung function development in children with versus without asthma, but the low numbers in our cohort may preclude us from detecting subtle differences with age.

It is a strength of our study that neonatal spirometry was performed using the raised volume rapid thoracoabdominal compression technique, providing volume-anchored measurements [10], which makes its results comparable with the forced volumes obtained by spirometry later in childhood. This is an important difference compared to previous studies of neonatal lung function in relation to asthma and lung function development during childhood, which used non-volume-anchored methods [12–14,28]. Assessment of tidal breathing patterns is one such method, which is less sensitive, has high intra-individual variability [29,30], and yields results not associated with volume-anchored measurements later in childhood [13].

One previous study of 95 children investigated airway reactivity from age 1 month throughout childhood, but used a non-volume-anchored assessment of histamine challenge that did not correlate with the FEV_1 -based histamine challenge results at school age, highlighting the same problem of using different approaches to determine neonatal lung function and lung function later in childhood [12,31,32].

The limitations of our study are the relatively small sample size, the limited ethnic variation, and the high-risk nature of the COPSAC₂₀₀₀ cohort, which hamper the generalizability of the findings. However, mothers without asthma might be reluctant to have their neonates investigated so thoroughly by lung function testing during sedation. Children of asthmatic mothers are a priori at higher risk of asthma and may experience more severe symptoms and have poorer lung function. However, the children not developing asthma also had asthmatic mothers, and we do not expect the relationship between symptoms and lung function development to be different due to asthma predisposition.

We observed z-score differences in lung function trajectories with magnitudes between 0.3 and 0.4 in children who developed asthma compared to children not developing asthma. This overall difference in lung function development is not in the abnormal range, which would be less than -2 z-scores, but it should be kept in mind that we excluded measurements obtained

during exacerbations, which may have diminished the overall difference between children with versus without asthma.

Interpretation

We found that airway obstruction and bronchial hyperreactivity related to asthma are fixed traits from age 1 month to age 13 years, without further deterioration from disease duration or improvement after symptom remission. These findings have important implications for our understanding of the underlying pathology, indicating that the symptomatic phases of childhood asthma are not causing the airway obstruction and bronchial hyperreactivity typical of asthma. Instead, we propose that airway obstruction and bronchial hyperreactivity are inherent traits that increase the risk of developing airway inflammation and asthmatic symptoms during childhood. This may explain the lack of effect from early intervention with ICS, which was unable to change the natural course of asthma with respect to both symptom burden and lung function development [2–6].

Our findings are consistent with a follow-up study of children diagnosed with asthma at age 9 years showing that individuals still experiencing symptoms at age 26 years had stable airway obstruction and bronchial reactivity without any progression from the assessments at age 9 years [33]. In addition, a 22-year follow-up of a cohort of 40-year-old adults also showed that reduced lung function at cohort inception remained stable over time [34]. Together with the present study, this suggests that airflow obstruction and bronchial hyperreactivity are inherent and stable deficits measurable already at age 1 month. Thus, even though lung function fluctuates with symptom load and anti-inflammatory treatment over time [35], the lung function trajectory is a static characteristic that might contribute to asthmatic symptoms, exaggerated bronchial reactivity, and intermittent airway obstruction.

Our findings contrast with those of a recent study investigating the development of lung function in children with asthma into adulthood [36], which suggested that the majority of children with asthma (75%) had an abnormal pattern of reduced growth and/or early decline in lung function, i.e., not a stable trait. This difference may be due to the fact that the study only enrolled 5- to 12-year-old children with uncontrolled chronic asthma (symptoms >2 days/week) and severe bronchial hyperreactivity at baseline (>20% drop in FEV₁ after methacholine challenge), resembling a minority population of the most severely affected children. Even more important, the participants with reduced growth in lung function already had significantly lower lung function at enrollment, suggesting that their lung function trajectories may be a static characteristic established in early childhood. In line with our findings, another recent study demonstrated increased airway resistance among children with persistent wheeze, which tracked from early to late childhood [37].

We suggest that the propensity to develop asthmatic symptoms due to intrinsic and extrinsic factors is increased by underlying inherent airflow obstruction and bronchial hyperreactivity, which are stable traits without progression due to triggering factors or increased symptom duration. On the other hand, we and other groups have shown that allergic sensitization after the age of 6 years, but not earlier in childhood, is associated with an increased risk of developing asthma [38,39], with different risks for different allergens [40]. This raises the possibility that development of allergy may have a deteriorating effect on the lung function trajectory. However, our data in general argue against such a hypothesis as development of airflow obstruction as measured by FEV_z, MMEF_z, and sRaw_z among children with asthma was independent of allergic sensitization at age 13 years, whilst there was a non-significant suggestion of increased airway reactivity among sensitized asthmatic individuals that was significant in the subgroup with sensitization to pollens, which were the most common allergens in our

cohort. These findings are in line with a study of 1,719 15-year-old participants in the German GINIplus birth cohort, which did not show any associations between sensitization and spirometric indices in children with asthma [41], but are in contrast to the American TENOR study of 1,261 children aged 6 to 17 years with severe or difficult-to-treat asthma, which showed an association between increased airflow limitation and higher IgE levels [42]. Importantly, both these studies were cross-sectional analyses; a longitudinal study showed that development of lung function from age 1 month until 11 years was unaffected by atopy [13].

Interestingly, we found that all children who fulfilled the diagnostic criteria for asthma compared to children never diagnosed with asthma had increased airway obstruction, i.e., reduced FEV_z and MMEF_z (age 1 month to 13 years) and increased sRaw_z (age 3 years to 13 years), irrespective of remission of symptoms or not. Thus, fixed airway obstruction was apparent in children with both a transient and a persistent phenotype and without differences in development of FEV_z, MMEF_z, and sRaw_z between the phenotypes, whereas bronchial reactivity to methacholine (PD_z) was more pronounced in children with persistent as opposed to transient symptoms. This suggests that airway reactivity in children with transient asthmatic symptoms declines with airway growth through childhood, whereas reduced forced flows and increased airway resistance are fixed traits among children with the transient phenotype, irrespective of airway growth. These findings align with findings from longitudinal lung function measurements from birth until age 16 years in the Tucson Children's Respiratory Study [43] and from age 3 years until age 11 years in the Manchester Asthma and Allergy Study [37], showing that children with persistent wheeze compared to never wheeze had increased and fixed airway obstruction, which was also apparent for children with early-transient viral-induced wheeze even though they outgrew their symptoms. Furthermore, a smaller Australian study from Perth also showed a suggestion of reduced lung function from birth until age 11 years in children with persistent wheeze, whereas no reduction of lung function was observed for other phenotypes of wheeze [12]. Overall, these studies together with our study suggest a common underlying lung function deficit driving early-transient, typically viral-induced symptoms and persistent symptoms, typically triggered by allergens, pollutants, or exercise.

Conclusion

In this study we observed that airway obstruction and increased bronchial reactivity associated with childhood asthma were established already at age 1 month, without further deterioration into early adolescence and without relation to asthma symptom duration or remission. This suggests that these chronic deficits are not the result of childhood asthma, but may instead contribute to asthma pathogenesis by increasing the risk of symptoms, exaggerated hyperreactivity, and airway obstruction, implying that preventive measures for improved lung health should focus on the pre- or perinatal period.

Supporting information

S1 STROBE Checklist. Checklist according to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines.

(DOC)

S1 Data. Data utilized for the analyses presented in the paper.

(XLSX)

S1 Text. Supplementary text, tables, and figures.

(DOCX)

Author Contributions

Conceptualization: Bo Lund Chawes, Jakob Stokholm, Klaus Bønnelykke, Hans Bisgaard.

Data curation: Henrik Wegener Hallas, Morten Arendt Rasmussen.

Formal analysis: Henrik Wegener Hallas, Morten Arendt Rasmussen.

Investigation: Henrik Wegener Hallas, Bo Lund Chawes, Lambang Arianto, Jakob Stokholm.

Project administration: Hans Bisgaard.

Supervision: Bo Lund Chawes, Jakob Stokholm, Klaus Bønnelykke, Hans Bisgaard.

Validation: Morten Arendt Rasmussen.

Writing – original draft: Henrik Wegener Hallas.

Writing – review & editing: Bo Lund Chawes, Morten Arendt Rasmussen, Lambang Arianto, Jakob Stokholm, Klaus Bønnelykke, Hans Bisgaard.

References

1. Al-Muhsen S, Johnson JR, Hamid Q. Remodeling in asthma. *J Allergy Clin Immunol*. 2011; 128(3):451–62. <https://doi.org/10.1016/j.jaci.2011.04.047> PMID: 21636119
2. Bisgaard H, Hermansen MN, Loland L, Halkjaer LB, Buchvald F. Intermittent inhaled corticosteroids in infants with episodic wheezing. *N Engl J Med*. 2006; 354(19):1998–2005. <https://doi.org/10.1056/NEJMoa054692> PMID: 16687712
3. Guilbert TW, Morgan WJ, Zeiger RS, Mauger DT, Boehmer SJ, Szeffler SJ, et al. Long-term inhaled corticosteroids in preschool children at high risk for asthma. *N Engl J Med*. 2006; 354(19):1985–97. <https://doi.org/10.1056/NEJMoa051378> PMID: 16687711
4. Murray CS, Woodcock A, Langley SJ, Morris J, Custovic A. Secondary prevention of asthma by the use of Inhaled Fluticasone propionate in Wheezy INfants (IFWIN): double-blind, randomised, controlled study. *Lancet*. 2006; 368(9537):754–62. [https://doi.org/10.1016/S0140-6736\(06\)69285-4](https://doi.org/10.1016/S0140-6736(06)69285-4) PMID: 16935686
5. Busse WW, Pedersen S, Pauwels RA, Tan WC, Chen Y-Z, Lamm CJ, et al. The Inhaled Steroid Treatment As Regular Therapy in Early Asthma (START) study 5-year follow-up: Effectiveness of early intervention with budesonide in mild persistent asthma. *J Allergy Clin Immunol*. 2008; 121(5):1167–74. <https://doi.org/10.1016/j.jaci.2008.02.029> PMID: 18405951
6. Szeffler S, Weiss S, Tonascia J, Adkinson NF, Bender, Cherniack, et al. Long-term effects of budesonide or nedocromil in children with asthma. *N Engl J Med*. 2000; 343(15):1054–63. <https://doi.org/10.1056/NEJM200010123431501> PMID: 11027739
7. Lezmi G, Gosset P, Deschildre A, Abou-Taam R, Mahut B, Beydon N, et al. Airway remodeling in pre-school children with severe recurrent wheeze. *Am J Respir Crit Care Med*. 2015; 192(2):164–71. <https://doi.org/10.1164/rccm.201411-1958OC> PMID: 25961111
8. Bisgaard H, Stokholm J, Chawes BL, Vissing NH, Bjarnadóttir E, Schoos A-MM, et al. Fish oil-derived fatty acids in pregnancy and wheeze and asthma in offspring. *N Engl J Med*. 2016; 375(26):2530–9. <https://doi.org/10.1056/NEJMoa1503734> PMID: 28029926
9. Chawes BL, Bønnelykke K, Stokholm J, Vissing NH, Bjarnadóttir E, Schoos A-MM, et al. Effect of vitamin D3 supplementation during pregnancy on risk of persistent wheeze in the offspring: a randomized clinical trial. *JAMA*. 2016; 315(4):353–61. <https://doi.org/10.1001/jama.2015.18318> PMID: 26813208
10. Loland L, Bisgaard H. Feasibility of repetitive lung function measurements by raised volume rapid thoracoabdominal compression during methacholine challenge in young infants. *Chest*. 2008; 133(1):115–22. <https://doi.org/10.1378/chest.07-1328> PMID: 17951623
11. Loland L, Buchvald FF, Halkjaer LB, Anhøj J, Hall GL, Persson T, et al. Sensitivity of bronchial responsiveness measurements in young infants. *Chest*. 2006; 129(3):669–75. <https://doi.org/10.1378/chest.129.3.669> PMID: 16537866
12. Turner SW, Palmer LJ, Rye PJ, Gibson NA, Judge PK, Cox M, et al. The relationship between infant airway function, childhood airway responsiveness, and asthma. *Am J Respir Crit Care Med*. 2004; 169(8):921–7. <https://doi.org/10.1164/rccm.200307-891OC> PMID: 14764431

13. Håland G, Carlsen KCL, Sandvik L, Devulapalli CS, Munthe-Kaas MC, Pettersen M, et al. Reduced lung function at birth and the risk of asthma at 10 years of age. *N Engl J Med*. 2006; 355(16):1682–9. <https://doi.org/10.1056/NEJMoa052885> PMID: 17050892
14. Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. Asthma and wheezing in the first six years of life. The Group Health Medical Associates. *N Engl J Med*. 1995; 332(3):133–8. <https://doi.org/10.1056/NEJM199501193320301> PMID: 7800004
15. Martinez FD, Morgan WJ, Wright AL, Holberg CJ, Taussig LM. Diminished lung function as a predisposing factor for wheezing respiratory illness in infants. *N Engl J Med*. 1988; 319(17):1112–7. <https://doi.org/10.1056/NEJM198810273191702> PMID: 3173442
16. Bisgaard H. The Copenhagen Prospective Study on Asthma in Childhood (COPSAC): design, rationale, and baseline data from a longitudinal birth cohort study. *Ann Allergy Asthma Immunol*. 2004; 93(4):381–9. [https://doi.org/10.1016/S1081-1206\(10\)61398-1](https://doi.org/10.1016/S1081-1206(10)61398-1) PMID: 15521375
17. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J*. 2005; 26(2):319–38. <https://doi.org/10.1183/09031936.05.00034805> PMID: 16055882
18. Bisgaard H, Klug B. Lung function measurement in awake young children. *Eur Respir J*. 1995; 8(12):2067–75. PMID: 8666102
19. Crapo RO, Casaburi R, Coates A, Enright PL, Hankinson J, Irvin CG, et al. Guidelines for methacholine and exercise challenge testing—1999. *Am J Respir Crit Care Med*. 2000; 161(1):309–29. <https://doi.org/10.1164/ajrccm.161.1.ats11-99> PMID: 10619836
20. Nielsen KG, Bisgaard H. Lung function response to cold air challenge in asthmatic and healthy children of 2–5 years of age. *Am J Respir Crit Care Med*. 2000; 161(6):1805–9. <https://doi.org/10.1164/ajrccm.161.6.9905098> PMID: 10852748
21. Beydon N, Davis SD, Lombardi E, Allen JL, Arets HGM, Aurora P, et al. An official American Thoracic Society/European Respiratory Society statement: pulmonary function testing in preschool children. *Am J Respir Crit Care Med*. 2007; 175(12):1304–45. <https://doi.org/10.1164/rccm.200605-642ST> PMID: 17545458
22. Bisgaard H, Hermansen MN, Buchvald F, Loland L, Halkjaer LB, Bønnelykke K, et al. Childhood asthma after bacterial colonization of the airway in neonates. *N Engl J Med*. 2007; 357(15):1487–95. <https://doi.org/10.1056/NEJMoa052632> PMID: 17928596
23. Schoos A-MM, Chawes BLK, Følsgaard NV, Samandari N, Bønnelykke K, Bisgaard H. Disagreement between skin prick test and specific IgE in young children. *Allergy*. 2015; 70(1):41–8. <https://doi.org/10.1111/all.12523> PMID: 25224528
24. Paganelli R, Ansotegui IJ, Sastre J, Lange CE, Roovers MH, de Groot H, et al. Specific IgE antibodies in the diagnosis of atopic disease. Clinical evaluation of a new in vitro test system, UniCAP, in six European allergy clinics. *Allergy*. 1998; 53(8):763–8. PMID: 9722225
25. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Softw*. 2015; 67(1):1–48.
26. Vissing NH, Jensen SM, Bisgaard H. Validity of information on atopic disease and other illness in young children reported by parents in a prospective birth cohort study. *BMC Med Res Methodol*. 2012; 12:160. <https://doi.org/10.1186/1471-2288-12-160> PMID: 23088330
27. Wanger J, Clausen JL, Coates A, Pedersen OF, Brusasco V, Burgos F, et al. Standardisation of the measurement of lung volumes. *Eur Respir J*. 2005; 26(3):511–22. <https://doi.org/10.1183/09031936.05.00035005> PMID: 16135736
28. Collins SA, Pike KC, Inskip HM, Godfrey KM, Roberts G, Holloway JW, et al. Validation of novel wheeze phenotypes using longitudinal airway function and atopic sensitisation data in the first 6 years of life: evidence from the Southampton Women's Survey. *Pediatr Pulmonol*. 2013; 48(7):683–92. <https://doi.org/10.1002/ppul.22766> PMID: 23401430
29. Ranganathan SC, Bush A, Dezateux C, Carr SB, Hoo A-F, Lum S, et al. Relative ability of full and partial forced expiratory maneuvers to identify diminished airway function in infants with cystic fibrosis. *Am J Respir Crit Care Med*. 2002; 166(10):1350–7. <https://doi.org/10.1164/rccm.2202041> PMID: 12421744
30. Ranganathan S, Hoo A, Lum S, Goetz I, Castle R, Stocks J. Exploring the relationship between forced maximal flow at functional residual capacity and parameters of forced expiration from raised lung volume in healthy infants. *Pediatr Pulmonol*. 2002; 33(6):419–28. <https://doi.org/10.1002/ppul.10086> PMID: 12001274
31. Palmer LJ, Rye PJ, Gibson NA, Burton PR, Landau LI, Lesouëf PN. Airway responsiveness in early infancy predicts asthma, lung function, and respiratory symptoms by school age. *Am J Respir Crit Care Med*. 2001; 163(1):37–42. <https://doi.org/10.1164/ajrccm.163.1.2005013> PMID: 11208623

32. Turner SW, Palmer LJ, Rye PJ, Gibson NA, Judge PK, Young S, et al. Infants with flow limitation at 4 weeks: outcome at 6 and 11 years. *Am J Respir Crit Care Med*. 2002; 165(9):1294–8. <https://doi.org/10.1164/rccm.200110-018OC> PMID: 11991882
33. Sears MR, Greene JM, Willan AR, Wiecek EM, Taylor DR, Flannery EM, et al. A longitudinal, population-based, cohort study of childhood asthma followed to adulthood. *N Engl J Med*. 2003; 349(15):1414–22. <https://doi.org/10.1056/NEJMoa022363> PMID: 14534334
34. Lange P, Celli B, Agustí A, Boje Jensen G, Divo M, Faner R, et al. Lung-function trajectories leading to chronic obstructive pulmonary disease. *N Engl J Med*. 2015; 373(2):111–22. <https://doi.org/10.1056/NEJMoa1411532> PMID: 26154786
35. Castro-Rodriguez JA, Rodrigo GJ. Efficacy of inhaled corticosteroids in infants and preschoolers with recurrent wheezing and asthma: a systematic review with meta-analysis. *Pediatrics*. 2009; 123(3):e519–25. <https://doi.org/10.1542/peds.2008-2867> PMID: 19254986
36. McGeachie MJ, Yates KP, Zhou X, Guo F, Sternberg AL, Van Natta ML, et al. Patterns of growth and decline in lung function in persistent childhood asthma. *N Engl J Med*. 2016; 374(19):1842–52. <https://doi.org/10.1056/NEJMoa1513737> PMID: 27168434
37. Belgrave DCM, Buchan I, Bishop C, Lowe L, Simpson A, Custovic A. Trajectories of lung function during childhood. *Am J Respir Crit Care Med*. 2014; 189(9):1101–9. <https://doi.org/10.1164/rccm.201309-1700OC> PMID: 24606581
38. Schoos A-MM, Chawes BL, Rasmussen MA, Bloch J, Bønnelykke K, Bisgaard H. Atopic endotype in childhood. *J Allergy Clin Immunol*. 2016; 137(3):844–51.e4. <https://doi.org/10.1016/j.jaci.2015.10.004> PMID: 26597163
39. Govaere E, Van Gysel D, Verhamme KMC, Doli E, De Baets F. The association of allergic symptoms with sensitization to inhalant allergens in childhood. *Pediatr Allergy Immunol*. 2009; 20(5):448–57. <https://doi.org/10.1111/j.1399-3038.2008.00805.x> PMID: 19175888
40. Sears MR, Herbison GP, Holdaway MD, Hewitt CJ, Flannery EM, Silva PA. The relative risks of sensitivity to grass pollen, house dust mite and cat dander in the development of childhood asthma. *Clin Exp Allergy*. 1989; 19(4):419–24. PMID: 2758355
41. Luzak A, Flexeder C, von Berg A, Berdel D, Klümper C, Koletzko S, et al. Relation of lung function and current inhalant allergen-specific immunoglobulin E concentrations in adolescents (GINIplus cohort). *Ann Allergy Asthma Immunol*. 2015; 115(3):183–90. <https://doi.org/10.1016/j.anai.2015.06.016> PMID: 26195441
42. Haselkorn T, Szefer SJ, Simons FER, Zeiger RS, Mink DR, Chipps BE, et al. Allergy, total serum immunoglobulin E, and airflow in children and adolescents in TENOR. *Pediatr Allergy Immunol*. 2010; 21(8):1157–65. <https://doi.org/10.1111/j.1399-3038.2010.01065.x> PMID: 20444153
43. Morgan WJ. Outcome of asthma and wheezing in the first 6 years of life: follow-up through adolescence. *Am J Respir Crit Care Med*. 2005; 172:1253–8. <https://doi.org/10.1164/rccm.200504-525OC> PMID: 16109980